

1 INTRODUCTION


The image of a human face contains various kinds of information relevant in the social context of recognition and communication. The human brain appears to be highly specialized to extract this information. We can identify a familiar person from a single view and we can make reliable judgements about age, sex and the facial expression of unfamiliar faces. To a great extent these abilities are invariant to changes of the orientation of a face in space.

Orientation itself is an attribute that humans seem to be able to derive from the image of a face. As human heads are approximately bilaterally symmetric, the orientation is well defined (e.g. with respect to the frontal view). Not only may knowledge about the orientation of a face play a role in achieving viewpoint invariance for recognition, but it may also contain important information about the direction of attention or movement.

What are the mechanisms underlying the ability to judge the orientation of a 3D object from a 2D image?

We approach this question by investigating a phenomenon that we discovered in the course of recent experiments dealing with the ability to generalize to new illuminations (Troje & Bühlhoff, 1996):

The perceived orientation of a head is significantly influenced by the illumination



Orientation (f_1) = +30 deg Illumination (l_1) = +30 deg
Orientation (f_2) = +30 deg Illumination (l_2) = -30 deg

In a series of four experiments, we will first demonstrate and quantify this phenomenon and then test for its dependence on the following parameters:

Mean orientation of the two faces (Exp. 1)

Magnitude of the illumination change (Exp. 2)

Visibility of gaze direction (Exp. 3)


Visibility of the occluding contour of the faces (Exp. 4)

2

Illumination induced apparent orientation shift (IAOS) was determined by measuring the physical orientation change necessary to compensate for it.

PROCEDURE:


- A single trial showed a sequence of two images (4 degs visual angle) of the same face separated by a distracting mask.
- Using a 2AFC paradigm subjects were asked to report the direction (right/left) of the perceived orientation change.
- Each trial was characterized by the following parameters:
 - f_m : mean orientation of the two images $f_m = (f_1 + f_2)/2$
 - Δf : physical orientation change $\Delta f = f_2 - f_1$
 - l_m : mean light source position $l_m = (l_1 + l_2)/2$
 - Δl : position change of the illumination $\Delta l = l_2 - l_1$



DESIGN:

- 5 blocks (80 trials each) per experiment: 4 conditions + 1 control
- Within each block only the physical orientation change was varied: 8 steps from -6 degs to 15 deg (with respect to the direction of Δl)
- The order of the blocks as well as the order of the trials within each block were randomized for each subject. 8 different subjects participated in each experiment.

EXAMPLE:




3

RESULTS


Exp. 1: Effect of the mean orientation

- black background
- $\Delta l = 60$ deg, $l_m = 0$ deg
- 4 conditions: $f_m = 0, 15, 30, 45$ deg
- control: $f_m = 0$ deg, $l_m = 0$ deg, $\Delta l = 0$ deg



IAOS increases significantly with mean orientation

$$F_{4,28} = 33.4, p < 0.01$$




Hypothetical predictors for face orientation:

- $\frac{w_r}{w_l}$
- $\frac{w_r}{h}$
- $\frac{A_r}{A_l}$


Exp. 2: Dependence on angle between light sources

- black background
- $f_m = 30$ deg, $l_m = 0$ deg
- 4 conditions: $\Delta l = 30, 60, 90, 120$ deg
- control: $f_m = 30$ deg, $l_m = 0$ deg, $\Delta l = 0$ deg



IAOS first increases with magnitude of the illumination change, but then saturates for $\Delta l > 60$ deg

$$F_{4,28} = 16.1, p < 0.01$$




$f_1 = +25.5$ degs, $l_1 = -30$ degs
 $f_2 = +34.5$ degs, $l_2 = +30$ degs

- perceptually equivalent orientations
- geometrical relations identical

Exp. 3: Influence of gaze direction


- black background
- constant albedo
- variables as in Exp. 1



IAOS similar to results in Exp. 1

Exp. 4: Importance of occluding contours

- beige background
- variables as in Exp. 1



No significant difference between treatments

IAOS reduced to values < 2 deg

5 REFERENCES

Troje, N. F. and Bühlhoff, H. H. (1996) *What is the basis for good performance to symmetric views of faces?* Paper presented at ARVO 1996 [Abstract published in Investigative Ophthalmology and Visual Science, 37 (3), S194.]

Troje, N. F. and Bühlhoff, H. H. (in press) *How is bilateral symmetry of human faces used for recognition of novel views?* Vision Research.

Koenderink, J. J., van Doorn, A. J. and Christou, C. (1996) *Shape constancy in pictorial relief.* Perception 25: 155-164.